



Datasheet for Lemon

Sunergist® EBR (24-Epibrassinolide 0.01% SL)

To lemon growers, two key indicators mean good return for fresh market: advanced harvest and more preferred size of fruits. Controlled growth at every key stage, with *Sunergist® EBR* will be possible. Key words: even flowering, even sizing, and even ripening.

Suggested Use Direction

- 1st spray - Pre-flowering (popcorn stage)
- 2nd spray - 7-10 days at petal fall with 6 fl. oz / 100 GPA
- 3rd spray - silver coin stage at 6 fl. oz / 100 GPA
- 4th spray - at early ripening stage is also suggested at 12 fl. oz / 100 GPA

Note: Nonionic surfactant (NIS) must be used.

Benefit to Growers

- Protect the pollination and increase fruit set.
- Enhancing uniformity of fruit growing.
- Improve the uniformity of fruit ripening and increase the sellable fruits for fresh market.

Please read label before use:

Crop	Amt of product in fl. oz./10 gals of water	Application timing
Citrus fruits (Crop Group 10-10) such as: Sweet orange, Lemon and Grapefruit	0.6 - 1.2	1 st . 5-10 days after first flower 2 nd . Apply when fruit size is 5 mm To increase doubles additional sprays maybe needed.

Our Distributor

California
Coastal - Helena Agri-Enterprises
Coachella - TBD

Arizona
Helena Agri-Enterprises