

Datasheet for Peanut

Sunergist® EBR (24-Epibrassinolide 0.01% SL)



Sunergist® EBR coincides with post emergence herbicide at 4-8 leaf stage will also enhance the healthy growth under herbicide shock. During R2-R3 fungicide time, coincides with Sunergist® EBR will improve the effective pegging and uniformity of pod development.

Suggested Use Direction

- 1st spray - 4-8 leaf stage, coincides as postemergence herbicides tan-mix with 0.6 fl. oz / 10 GPA
- 2nd spray - during R2 to R3, coincides as fungicides tank mix with 0.6 fl. oz / 10 GPA (repeat with more fungicide applications)

Note: Nonionic surfactant (NIS) must be used

Benefit to Growers

- Enhancing the uniform growth and alleviate toxicities from herbicide application
- Help for reduce the plant height and effective pegging.
- Sunergist® EBR will improve the uniformity of kernel development.

Please read label before use:

Crop	Amt of product in fl. oz./10 gals of water	Application timing
Peanut	0.6 - 1.2	1 st . At initial blooming 2 nd . At early R2 (tank mix with fungicide) 3 rd . At early R3 (tank mix with fungicide)

Our Distributor

US Southeast
CNI member distributors
Helena Agri-Enterprises