



Datasheet for Table Grape

Sunergist® EBR (24-Epibrassinolide 0.01% SL)

To table grape growers, gaining uniformity ripening with high quality fruits would be ultimate goal. All facts affect the uniformity of ripening, such as even flowering and sizing treated by Sunergist® EBR. As well during ripening stage.

Suggested Use Direction

- 1st spray - white bud stage with 12 fl. oz / 100 GPA
- 2nd spray - sizing stage @ 8-10mm with 12 fl. oz / 100 GPA, tank-mixing with GA3 application
- Ripen spray - 20-30% of berry softening stage with 51.2 - 64 fl. oz / 100 GPA

Note: Nonionic surfactant (NIS) must be used

Benefit to Growers

- Enhancing flower bud development for even flowering
- Ensure even sizing
- Promoting the uniformity of fruit ripening and help for coloration of red variety
- Cytokinin (as kinetin) are recommended to use coincide with Sunergist® EBR in all applications for increasing cell divisions

Please read label before use:

Crop	Amt of product in fl. oz./10 gals of water	Application timing
Table Grape	1.2 - 6.4	1 st . At bud break 2 nd . At small fruit sizing (Tank-mix with GA3) 3 rd . At berry softening (Enhance ripening)

Our Distributor

California

Coachella - TBD

Kern County - Helena Agri-Enterprises

San Joaquin - Helena Agri-Enterprises